THE FREMONT-RIDEOUT HEALTH GROUP

ANAPHYLACTIC SHOCK ORDERS

IMPRINT PATIENT PLATE WITHIN THIS BOX

FAX ORDERS TO PHARMACY ASAP

AFTER INITIAL SHOCK ORDERS

Goals: MAP ≥ 70

U/O ≥ 1/2 mL/kg/hour

Good peripheral perfusion with capillary refill ≤ 3 seconds

Mixed venous SvO_2 ≥ 60; Central Venous SvO_2 ≥ 70

CI ≥ 3.0

1. Epinephrine 1:10,000 (0.1 mg/mL) -- 0.5 mL IVP Q 5 minutes PRN until SBP > 90 followed by epinephrine 5 mg/250 mL NS infusion at initial rate of _____, titrate to maintain SBP > 90 or MAP > 60

2. 500 mL LR, IV bolus over 10 minutes

3. Nebulized racemic epinephrine

4. Methylprednisolone (Solu-Medrol) 125 mg IVP STAT

5. Diphenhydramine (Benadryl) 1 mg/kg IVP STAT

6. Famotidine (Pepcid) 80 mg IVP STAT

7. If on norepinephrine (Levophed) or epinephrine drip when MAP ≥ 70, add dobutamine 2 mcg/kg/minute

8. Continue **FLUID CHALLENGE ORDERS** to maintain goals

Unit Clerk: Date: Time: Noted by RN: Date: Time: